

Diabetes mellitus in Cats

What is Diabetes Mellitus?

Diabetes mellitus is also called "sugar diabetes," and occurs when there are high glucose (sugar) levels in the blood and urine. This can happen either as a result of the failure of the pancreas to make insulin (type I diabetes) or the failure of the cells in the body to respond to insulin (type II diabetes). The more common form, type I, or insulin dependent diabetes mellitus (IDDM), usually requires insulin injections for treatment. Type II diabetics may not require insulin for treatment initially, but will often eventually develop the need for insulin seen in type I diabetics. Type II diabetes is also called non-insulin dependent diabetes mellitus (NIDDM). There are no reliable tests to tell type I from type II diabetes – response to insulin over time tells which type a cat has. Diabetes mellitus is one of the most common endocrine (hormonal) diseases in cats.

Transient diabetes mellitus can occur in cats. Unlike permanently diabetic cats, transient diabetics may require treatment with insulin only for a brief period of time. Transient diabetes can be seen in overweight cats, cats with pancreatitis or other diseases, or cats on medications that interfere with insulin activity. Some transient diabetics require insulin for a short period of time, then never again. Others need insulin from time to time at various periods during their life. Still others eventually become permanent diabetics, and ultimately need lifelong daily insulin therapy.

What are the Symptoms of Diabetes?

Diabetes mellitus often is suspected because of the presence of the classic signs of increased thirst, more frequent urination, and weight loss often despite a greater appetite. Most cats are older than 10 years of age at the time of diagnosis. However, diabetes mellitus can be diagnosed at any age. Male cats are diagnosed more frequently than females, and all breeds can be affected. Owners may notice that they have to change the litter box more frequently because it is wet all the time. Some cats will begin to void large amounts of urine in places other than the litter box. This may mistakenly be perceived as a behavioral problem in some cats.

Over time, diabetes can lead to:

- delayed healing and chronic skin conditions – this is why we want to do bloodwork on animals that come in "just for skin problems." Diabetic skin problems can look just like allergic skin problems.
- walking on the hocks, or hyperextended wrists.
- Infection – especially in the urinary tract.

If left untreated, diabetes can progress further to weakness, dehydration, depression, diabetic coma, and even possibly death.

What Causes Diabetes?

Development of diabetes mellitus may be influenced by the presence of complicating factors such as obesity, concurrent disease, or medications that interfere with insulin's activity. Pancreatitis (inflammation of the pancreas), whether acute or chronic, is known to lead to

diabetes, as the pancreas makes the insulin needed to control blood sugar. Medications that can cause cats with borderline insulin production to become dependent on insulin include anti-inflammatory steroids (prednisone, dexamethasone, DepoMedrol, etc.) and progesterones such as MegAce.

How is Diabetes Diagnosed?

Diabetes is sometimes diagnosed as the result of routine blood test – there is excessive glucose in the blood and in the urine. Because stress can cause significant elevations of blood glucose levels in cats, and in some circumstances, glucose can be seen in the urine of stressed cats, the measurement of fructosamine may help distinguish stress-induced changes in blood and urine sugar levels from true diabetes mellitus. In stressed cats, serum fructosamine concentrations are usually normal, but they are elevated in diabetic cats. Fructosamine indicates an average of blood glucose levels over the previous 10-14 days. Ketones are another substance that can appear in the blood and urine of diabetic cats, but they are generally not seen due to stress.

Ideally, the following tests should be done on every new diabetic cat, prior to pursuing long term management, to assess the condition of the diabetic cat, and to discover at the outset concurrent medical conditions that might change long term prognosis, or hamper insulin regulation.

- *CBC (complete blood count)* – high white count might indicate infection that needs to be isolated and treated. Anemia, if present, should also be treated.
- *Biochemistry profile* – to look for liver or pancreas problems that commonly occur in unregulated new diabetic cats, and may need special treatment, as well as to look for problems such as kidney disease which are common in older cats. The profile must include phosphorus in sick diabetic cats, as phosphorus can become dangerously low in new diabetics.
- *Electrolytes* – low potassium is common in new diabetics, and can be life threatening in diabetics with ketoacidosis.
- *fPLI or TLI* – many diabetic cats have chronic and/or acute pancreatitis which can be detected by these tests, and must be treated if the cat is to do well. A few with long standing chronic pancreatitis can develop enzyme deficiencies which must be treated by adding enzymes to each meal, so that their body can put to use the nutrients in food.
- *Thyroid test* - should be checked in geriatric cats to rule out concurrent hyperthyroidism (overactive thyroid), which can hamper efforts to get diabetic cats regulated.
- *Urinalysis and urine culture* - necessary to rule out ketoacidosis and urinary tract infection. A significant proportion of new diabetics have a urinary tract infection.
- *Chest x-rays and abdominal ultrasound* - may also be recommended, depending on the cat's overall condition, to assess for concurrent problems which could change prognosis and deter an owner from pursuing long term management of a diabetic.
- *FeLV antigen and FIV antibody tests*- owners should know the feline leukemia and FIV (cat AIDs) status of their cat prior to pursuing lifelong treatment.

Which Insulin should I use?

The commonly used insulins are listed below, listed from shortest to longest acting:

- “R” for regular – the shortest acting – usually used only for cats that are ill in the hospital.
- “N” for NPH – rarely used for cats, because it is usually too short acting. Most cats eliminate it in just a few hours.
- “L” for Lente – trade name Vetsulin; available only from veterinarians and veterinary pharmacies. Recent problems with manufacturing have made use of this insulin problematic, and soon it will no longer be available.
- “PZI” for Protamine Zinc Insulin – available only from veterinarians and veterinary pharmacies. A good insulin for cats, but can be very expensive.
- Lantus of glargine insulin – the longest acting, and usually the best for cats, as cats tend to metabolize insulin quickly.

The manufacturer of Lantus recommends that the bottle be discarded 28 days after opening. However, independent studies show that Lantus is effective for at least 6 months after opening, assuming that the expiration date has not passed. If the insulin becomes discolored or cloudy or changes appearance in any way, it should be discarded.

IMPORTANT!!!! There are two different concentrations of insulin, called U-100 and U-40. Each type of insulin has its corresponding calibrated syringes, also called U-100 and U-40. **You MUST know which your cats take, so you use the right syringes.** U-100 insulin contains 100 units per cc (Lantus and NPH), and U-40 insulin contains only 40 units insulin per cc (Vetsulin, PZI and ProZinc). Giving Lantus or NPH (U-100) insulin using the wrong (U-40) syringes will result in a 250% overdose. Giving Vetsulin, PZI or ProZinc (U-40) insulin using U-100 syringes will result in giving only 40% of the insulin needed.

- U-40 syringes usually have red caps – for Vetsulin, PZI and ProZinc.
- U-100 syringes usually have orange caps – for NPH and Lantus.

Cats are usually on smallish dosages of insulin, so the small-capacity (3/10 or 1/2 cc) syringes work well.

Insulin is usually given to cats every twelve hours. Ideally, insulin should be given no closer together than 10 hours, and never farther apart than 14 hours.

What Diet should Diabetic Cats Eat?

The proper diet is EXTREMELY IMPORTANT to getting your diabetic cats regulated. As a general rule, semi-moist foods are avoided, because they are high in sugars and carbohydrates. High protein (>40% of calories) and low carbohydrates (<10% of calories) are indicated, as some cats will have lower insulin requirement, or even go off insulin altogether if fed high protein/low carbohydrate diets. There are many canned diets on the market which fit the bill, but only two dry diets, to my knowledge – Innova EVO and Wellness Core. Purina DM and Hill’s Prescription Diet MD have sufficient protein, but are 15% carbohydrates, which is a big improvement over the 25-35% carbs in most commercial dry feline diets, but not ideal for diabetics.

Especially for overweight cats, adding high fiber to the diet may be helpful in encouraging weight loss and in controlling fluctuations in the serum glucose concentrations. In the past, commercially prepared high fiber diets such as Hills Prescription Diet R/D and W/D were recommended for diabetic cats. However, these diets are too high in carbohydrates and too low in protein for diabetic cats. If fiber is to be increased in the diet, psyllium (1/4 to 1/2 tsp per 3 ounces of canned food) or a small spoon of pumpkin can be added to a high

protein diet. Most diabetic cats do well on high protein/low carb diet, and don't need increased fiber to do well.

If a cat refuses to eat the proper diet, then other diets should be used, and insulin should be given accordingly. Frequency of feeding is usually dictated by the cat's normal dietary behavior. Cats that eat all day long are probably best fed small amounts many times per day, while cats that eat voraciously but infrequently may do better with meal feedings.

Phases of Treatment

Now that we have covered diet and insulin, there are three phases in treatment of diabetics: regulation, maintenance, and remission.

Regulation. It can take anywhere from 1-4 months to find the proper dose of insulin for your cat – this process is called "regulation." We usually keep a new diabetic for 1-2 days to monitor response to the first few doses of insulin, then send your pet home on a relatively low dose of insulin, and recheck them in 1-2 weeks. Unless you can do an entire glucose curve at home (check blood sugar every 1-2 hours throughout the day), your cat will need to stay at the clinic for 8-24 hours for these "regulation rechecks" in order to complete the glucose curve. Insulin dose is changed every 1-2 weeks, according to each glucose curve, until the ideal insulin dose is reached. If your cat is ill enough to be hospitalized at the time of diagnosis, we may hospitalize more often during the regulation period so that we can arrive at the proper dose more quickly.

The goal of regulation is to keep the lowest blood glucose number in the curve above 80 mg/dl, and the highest number below 200 mg/dl. Sometimes we have to settle for a higher peak, but we NEVER want blood glucose to go below 80 if we can help it, and we'd like to have the lowest blood glucose level below 120.

Once your cat is regulated, keep in mind that the proper insulin dose for your cat is subject to change. Coming in for scheduled rechecks and monitoring for signs of low blood sugar (see below) as well as PU-PD can detect need to change insulin dose.

Maintenance. When the cat is regulated to our satisfaction, you get a big chunk of your life back. Sometimes, it is no longer necessary to do glucose curve for very long periods of time (months to years). These are the regular checks we recommend for diabetic cats:

- **3-6 times a year (every 2-4 months)** - Short visit for weight and fructosamine (blood test). Healthier diabetics come less often, and fragile diabetics come more often. Start with every 2 months, and go from there.
- **Twice a year (every 6 months)** - Full exam including blood pressure, Complete Blood Count (CBC – to look for anemia or infection), general health profile (to check kidney, liver, etc), urinalysis, and urine culture (to check for bladder infection).
- **Once a year** - Thyroid testing yearly if your cat is older than 10 years old, or sooner if weight loss becomes a problem. Feline leukemia and FIV test if your cat goes outside.
- **Regular dental cleaning** as needed to prevent gum and tooth infections, which can dysregulate diabetic cats
- Any other tests or treatments indicated for your individual cat

Remission. Some very lucky cats actually have periods where they are not on medications (including insulin) and you can't tell they have diabetes. Remissions can be temporary or permanent. They can be short or long.

How Can I Monitor My Diabetic Cat?

PU-PD. The simplest way to monitor your cat is to pay attention to how much water he/she is drinking, and how much urine he/she is producing (watch the litter box). Increase in either should trigger you to check your cat's blood sugar, and visit with your vet about what to do next.

Fructosamine. The vets can do blood tests (fructosamine) that show how well the diabetes has been controlled over the past 10-14 days. If your cat who was previously well regulated is showing signs of poor regulation (weight loss, PU-PD, increased appetite, illness, etc), it's usually time to see the vet to check fructosamine. If fructosamine is normal, you assume the diabetes has been well controlled for the past 2 weeks, and the insulin dose is appropriate for your pet at that time. If fructosamine is elevated, then we know that insulin dose needs to be adjusted -- its' time for a glucose curve. Fructosamine can be elevated if the insulin dose is too low, or if the insulin dose is too high.

Home blood sugar testing. Checking blood sugar at home can make caring for your diabetic cat much easier and more effective. It might sound intimidating at first, but almost all owners can learn to check blood sugars at home. If you can check blood sugars at home, you and your vet can manage insulin treatments while keeping visits to the clinic to a minimum. As well, any time a cat is stress; blood sugar levels can go up, so blood sugar testing at the clinic likely is not as accurate as testing at home.

Home Blood Testing involves pricking the skin somewhere to obtain a drop of blood. Home blood testing can range from the occasional check to confirm your suspicion of hypoglycemia, to taking full control of regulating the cat's diabetes and doing glucose curves.

Glucometers made for human diabetics are used, and provide sufficient accuracy for optimal blood glucose control. These meters are very accurate in the normal ranges, less so as the glucose levels approach extremes. This doesn't matter, since all you need to know is that an extreme has been reached -- the extent of the extreme is less important than the fact that the glucose is far from where you want it.

In cats, the ear is generally the best site for blood testing. Although sensitive to touch, it is a relatively insensitive area to prick (lancets cause less pain than a mosquito bite). A paw pad can be used, but since the paws are on the floor and in the litter box, more care has to be taken about keeping the little puncture wound clean to avoid infection. Some are able to learn to use a needle to draw a few drops of blood from a leg vein, but this isn't for everyone.

There are several detailed handouts on how to do ear sticks so I won't repeat any of that here. Each person has to develop their own way of getting the sample from their own cat. If you are having trouble, let us know -- we've got lots of helpful hints to share.

Helpful hints for home blood glucose testing:

- Try to get a glucometer which uses a tiny volume of blood (One Touch Ultra) -- it is MUCH easier to use.
- Test strips are extremely sensitive to humidity and substances that might be in the air, so keep the vial on the test strip lid on tight.

- Never make insulin changes based on a single blood test reading. If single test readings are too high or too low, we need to do an entire curve to tell how to adjust the dose.
- Test strips are easily contaminated -- wash your hands in case you wind up accidentally touching it, and don't use it if it dropped on a dirty surface.
- Make sure the number on your test strips matched the number set on your glucometer (see your glucometer owners manual for more details on this, or ask us to show you).

Urine Dipsticks. Some use urine glucose testing for monitoring, but we don't, because it does not tend to be very accurate. We get better results with blood testing.

Hypoglycemia (low blood sugar)

Low blood sugar (hypoglycemia), if mild, causes little more than discomfort and hunger. But if BG level falls too far, you might see strange behavior, distress, or twitching. If BG falls dangerously low, problems can advance to severe distress and possibly seizures, coma, and even death in very severe cases. Some animals are very sensitive to even mild hypoglycemia; others give no obvious clues even when the levels are getting into what is generally thought to be dangerous territory. With time, you will learn how your cat responds to low blood sugar.

Mild Hypoglycemia. Signs of mild hypoglycemia (glucose 60-80 mg/dl):

- more quiet than usual
- excessive hunger
- sometimes a little bit of weakness or incoordination.

Mild hypoglycemia can usually be easily managed with an extra feeding of the regular diet, or a high protein canned food, meat or cheese. Check the food dish to make sure your cat has been eating normally. Check the litter box for signs of diarrhea, and check around the house for vomit, in case he ate and brought it back up.

If you find no evidence that a temporary shortage of food was the cause of low blood sugar, the next insulin dose should be skipped, and then the dose should be reduce by about 10-15% when insulin is given again. Two weeks after decreasing the dose, you should take your cat in to have fructosamine checked, or do a glucose curve at home or at the clinic, to see if the new dose is appropriate. If the cat is doing exceptionally well, it may be sufficient to just check a fructosamine. If there are any doubts that the dose is correct, a curve is better.

Moderate Hypoglycemia. Signs of moderate hypoglycemia (glucose 45-60 mg/dl)

- Twitching
- Weakness or incoordination
- Staggering, circling, huddling, trembling
- Disoriented, howling
- unable to see or correctly look in the direction of a noise

For moderate hypoglycemia, immediately offer your cat 1-3 cc of corn syrup (make sure you keep a syringe on hand – we can give you one). As soon as you see that his coordination is improving, give him a small snack of high protein cat food, cheese, meat or cat treats. In 30-60 minutes after initial improvement, offer a small serving of his regular food. If your cat is not back to normal within 1-2 hours, you need to make a visit to your

vet. Skip the next dose of insulin, and then decrease insulin dose by 20-25% when insulin injections are resumed. Do a fructosamine or glucose curve 1-2 weeks after adjusting the dose

Other high sugar liquids that can be used in place of corn syrup include dextrose solution (from your vet), honey, pancake syrup, even table sugar dissolved in water. HINT: keep the top of the corn syrup bottle clean so the cap doesn't get glued on -- if it's stuck shut while your cat is having an emergency your own anxiety levels may go through the roof.

Should he go to the vet, or be left to recover? That depends on whether you can be there to watch for a few hours, and your own confidence levels.

Severe Hypoglycemia. Signs of severe hypoglycemia (glucose less than 45 mg/dl):

- collapse
- blindness
- seizures
- howling
- comatose

In the case of severe hypoglycemia, put corn syrup on the cat's gums (be careful not to get bitten if your cat is disoriented or having seizures), and THEN call the vet on call. If you are having trouble with the corn syrup, just hold your cat firmly by the scruff (you won't hurt him by doing this), and squirt it on his tongue and gums. Don't try to restrain the seizures, but make sure that your cat is in an open space if having seizures, so he does not injure himself. As soon as you can, get your cat to a veterinarian – this type of hypoglycemia can be potentially life threatening. Your vet will make recommendations for insulin administration.

What Health Challenges Might I Expect in my Diabetic Cat (Prognosis)?

The outlook for diabetic cats depends on a number of factors. The owner's commitment to treatment is a key point, as treatment is seldom easy and is usually life-long. Cats diagnosed with diabetes mellitus at a younger age that are able to be controlled with relative ease can live for many years with a good quality of life. Cats that develop diabetes mellitus along with other diseases may also have a worse prognosis. One study that looked at the length of survival following the diagnosis of diabetes mellitus in cats found that the cats in the group examined live an average of two years after diagnosis. Many of our patients live much longer than that, once diagnosed with diabetes. Proper home care, regular veterinary evaluation, and most importantly excellent client-veterinary communication are vital to the successful treatment of diabetes mellitus in cats.

Cats are pretty tough creatures, and seem to resist the ravages of diabetes better than, say, dogs, who can go blind from cataracts in a matter of months if their blood glucose isn't controlled. Their insulin needs might be pretty constant, or they might change. If your cat is well regulated, he should remain in relatively good health. If you see any of the problems listed at the top of this handout as signs of diabetes, consider bringing him in for a check-up to see if he really is well regulated. If you monitor blood sugar at home, any loss of regulation that lasts more than a few days, or inability to achieve regulation, warrants a thorough checkup to see if something else is going on.

Keep a close eye on diabetics with early kidney failure -- if it worsens, the symptoms are increased water drinking and urination, the same as for diabetes out of control. You will need to have your vet check kidney functions periodically (by blood test).

Helpful Hints for Diabetics

If your cat wears a collar, make sure the tag announces that he is diabetic and includes your phone number and/or the vet's.

Corticosteroids such as prednisone and dexamethasone and hydrocortisone can potentially dysregulate diabetics, and large doses have actually induced diabetes in many animals. Unless your cat has a condition such as inflammatory bowel disease which necessitates use of prednisone, corticosteroids should be avoided in diabetics.

Any illness that lasts more than a couple of days will often necessitate an insulin change, at least temporarily. Sick cats eat less, move around less, and use less glucose.

If your cat ever needs anesthesia for surgery, dental cleaning, etc., DO NOT WITHOLD WATER OVERNIGHT, as many vets instruct owners to do with healthy cats. Take food up several hours after the evening insulin dose. No food in the morning, and take the water up then. Administer half of the regular insulin dose that morning. Give half the regular insulin dose the night after anesthesia, and then go back to the regular routine the next day.

References:

1. <http://felinediabetes.com>
2. <http://vetcentric.com>